



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## DIOPHANTINE ANALYSIS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

### SOLUTIONS TO PROBLEMS.

6. Proposed by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Find three whole numbers the sum of any two of which is a cube.

II. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

Let  $x$ ,  $y$ , and  $z$  = the numbers. Then  $x+y=a^3$ ,  $x+z=b^3$ , and  $y+z=c^3$ ;  
in which  $c > b > a$ . Hence  $x+y+z = \frac{a^3+b^3+c^3}{2}$ .

$$\therefore x = \frac{a^3+b^3+c^3}{2} - c^3; y = \frac{a^3+b^3+c^3}{2} - b^3; z = \frac{a^3+b^3+c^3}{2} - a^3.$$

In order that each of the numbers be integral, *either* each of the cubes must be even, *or* two of the cubes must be odd and one even. Also, in order that each of the numbers be positive,  $a^3+b^3 > c^3$ .

The simple rule for finding the three numbers is as follows: Take three cubes fulfilling the above two conditions, and from half their sum subtract each cube separately; the remainders will be the three numbers required.

It is evident that many sets of numbers be obtained. We will illustrate by the following consecutive cubes: 1, (2) 8, (3) 27, (4) 64, (5) 125, (6) 216, (7) 343, (8) 512, (9) 729, (10) 1000, (11) 1331, (12) 1728, (13) 2197, (14) 2744, (15) 3375, (16) 4096, (17) 4913, (18) 5832.

The first three cubes answering the above conditions are 343, 512, and 729.  $\frac{1}{2}$  their sum is 792;  $x=792-729=63$ ;  $y=792-512=286$ ;  $z=792-343=449$ . The next three cubes are  $(7)^3$ ,  $(9)^3$ , and  $(10)^3$ . The first three *even* cubes are  $(12)^3$ ,  $(14)^3$ , and  $(19)^3$ .

9. Proposed by ISAAC L. BEVERAGE, Monterey, Virginia.

It is required to find three numbers the sum of whose 4th powers is a square.

I. Solution by R. J. ADCOOK, Larchland, Warren County, Illinois.

If  $u = x^2 + y^2 - z^2$ , then  $u^2 = x^4 + y^4 + z^4 + 2x^2y^2 - 2x^2z^2 - 2y^2z^2$ , in which if  $2x^2y^2 - 2x^2z^2 - 2y^2z^2 = 0$ ,  $z^2 = \frac{x^2y^2}{x^2+y^2}$ , and  $u = x^2 + y^2 - \frac{x^2y^2}{x^2+y^2}$   
 $= \frac{x^4 + y^4 + x^2y^2}{x^2+y^2}$ ;  $u^2 = x^4 + y^4 + z^4 = x^4 + y^4 + \frac{x^4y^4}{(x^2+y^2)^2} = \left( \frac{x^4 + y^4 + x^2y^2}{x^2+y^2} \right)^2$ .

$\therefore x^4(x^2+y^2)^2 + y^4(x^2+y^2)^2 + x^4y^4 = (x^4 + y^4 + x^2y^2)^2$ , which is a general equation for the sum of the 4th powers of three quantities = a square, when  $x^2 + y^2 =$  a tation square. Making  $x=3, y=4$ ,  $x^2 + y^2 = 5^2$ , and  $3^4 \times 5^4 + 4^4 \times 5^4 + 3^4 \times 4^4 = 15^4 + 20^4 + 12^4 = (3^4 + 4^4 + 12^4)^2 = 481^2 = 231361$ .